

The more precise determination of hadronic contribution to muonic $(g - 2)$ factor and to $\alpha(M_z^2)$.

B.V.Geshkenbein *

Institute of Theoretical and Experimental Physics,
B.Chermushkinskaya 25, 117218 Moscow, Russia

Abstract

The hadronic vacuum-polarization contribution to the muon anomalous magnetic moment $a_\mu(hadr.)$ and to the value of the electromagnetic coupling constant at $q^2 = M_z^2$ are determined more precisely in connection with a new more exact value of the electronic width of ρ -meson. The QCD model with infinite number of vector mesons is used. The more precisely determined results are: $a_\mu(hadr.) = 678(7) \cdot 10^{-10}$, $\delta\alpha_{hadr.}(M_z^2) = 0.02786(6)$.

The purpose of this work is to determine more precisely the contribution of strong interactions to the value of $a_\mu(hadr.)$ and $\delta\alpha_{hadr.}(M_z^2)$ obtained in papers [1,2]. There are two reasons to write this letter.

1. A new more exact value of the electronic width of ρ -meson appeared [3]:

$$\Gamma_0^{ee} = 6.85 \pm 0.11 keV \quad (1)$$

instead of the value $\Gamma_0^{ee} = 6.77 \pm 0.32 keV$ used in [1] and of the value $\Gamma_0^{ee} = 6.72 \pm 10 keV$ used in [2], obtained by the author in paper [4], from the analysis of the old experiments on the measurement of the pion electromagnetic form factor.

2. Function $R(s)$ is calculated by new formula obtained in paper [5] where analyticity requirement of the QCD polarization operators were combined with the renormalization group. Function $R(s)$ for 3 flavours has the form

$$R(s) = \frac{3}{2}(1 + r(s)) \quad (2)$$

Function $r(s)$ is calculated [5] using the renormgroup and the requirement of absence of nonphysical singularities. All the formulae for calculation of the function $r(s)$ are given in paper [5]. The QCD model with and infinite number of vector mesons developed in papers [4,6,7] is the basis for the calculations. This model allows one to satisfy the requirements of the Wilson operator expansion and the nonperturbative effects [8] are included into it automatically. Such a model is very useful for calculation of the integrals of the function

*E-mail address: geshken@heron.itep.ru

$$R(s) = \frac{\sigma(e^+e^- \rightarrow \text{hadrons})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)} \quad (3)$$

Function $R(s)$ in this model has the form

$$R(s) = R_{ud}^{I=1}(s) + R_{ud}^{I=0}(s) + R_s(s) + R_c(s) + R_b(s) \quad (4)$$

Here $R_{ud}^{I=1}$ and $R_{ud}^{I=0}$ are the contributions of u and d quarks in the state with isotopic spin $I = 1$ (ρ family) and $I = 0$ (ω family) and $R_s(s)$, $R_c(s)$, $R_b(s)$ are the contributions of s (φ family), c (J/ψ family) and b (Υ family) quarks, respectively. The narrow resonances approximation is used. If the total width of the resonances $\Gamma_k \ll M_k$ (M_k is the k -th resonance mass) for all k , the narrow resonances approximation is valid. If $M_k \Gamma_k \gg M_k^2 - M_{k-1}^2$, then beginning from the k -th resonance function $R(s)$ will be described by a smooth curve and all the formulae of the model will be valid [9]. To calculate the integral of $R(s)$ in the QCD model with infinite number of vector mesons one should only know a few number of masses and electronic widths of lowlying resonances in each family [1,2]. Comparing with [1,2] only $R_{ud}^{I=1}(s)$ had been corrected. The changes in the contributions of the rest families are very small. All the formulae of papers [1,2] except for the formulae for $R_{ud}^{I=1}(s)$ are unchanged.

The results of the calculations for hadronic contribution of the muon ($g - 2$) factor are the following:

$$a_\mu(\text{had.}) = \frac{\alpha^2}{3\pi^2} \int_{4m_\pi^2}^{\infty} ds K(s) R(s)/s = 678(7) \cdot 10^{-10} \quad (5)$$

where

$$K(s) = x^2(1 - x^2/2) + (1 + x^2)(1 + x^{-2}) \left[\ln(1 + x) - x + x^2/2 \right] + \frac{1+x}{1-x} x^2 \ln x; \quad x = \frac{1 - (1 - 4m_\mu^2/s)^{1/2}}{1 + (1 - 4m_\mu^2/s)^{1/2}} \quad (6)$$

The result (5) should be compared with the recent, the most exact calculations $\alpha_\mu(\text{had.})$ obtained by integrating formula (5) with the experimental cross section of e^+e^- annihilation into hadrons [10,11].

$$a_\mu(\text{had.}) = (684.7 \pm 7) \cdot 10^{-10} \quad [10] \quad (7)$$

$$a_\mu(\text{had.}) = (681.1 \pm 6.3) \cdot 10^{-10} \quad [11] \quad (8)$$

For hadronic contribution into $\alpha(M_z)$ we have obtained

$$\delta\alpha_{\text{had.}} = \frac{\alpha M_z^2}{3\pi} P \int_{4m_\pi^2}^{\infty} \frac{R(s) ds}{(M_z^2 - s)s} = 0.02786(6) \quad (9)$$

The result (8) can be compared with the results of papers [12-22]:

$$\delta\alpha_{\text{had.}} = 0.02744(36) \quad [12]$$

$$\delta\alpha_{\text{had.}} = 0.02803(65) \quad [13]$$

$$\delta\alpha_{hadr.} = 0.02780(6) \quad [2]$$

$$\delta\alpha_{hadr.} = 0.0280(7) \quad [14]$$

$$\delta\alpha_{hadr.} = 0.02754(46) \quad [15]$$

$$\delta\alpha_{hadr.} = 0.02784(22) \quad [16]$$

$$\delta\alpha_{hadr.} = 0.02778(16) \quad [17]$$

$$\delta\alpha_{hadr.} = 0.02779(20) \quad [18]$$

$$\delta\alpha_{hadr.} = 0.02770(15) \quad [19]$$

$$\delta\alpha_{hadr.} = 0.02787(32) \quad [20]$$

$$\delta\alpha_{hadr.} = 0.02778(24) \quad [21]$$

$$\delta\alpha_{hadr.} = 0.02741(19) \quad [22]$$

obtained by integrating over eq.(8) with experimental cross section for e^+e^- annihilation into hadrons.

Acknowledgements

It was possible for us to do the research described in this publication in part by the Award No.RP2-2247 of the US Civilian Research and Development Foundation for the Independent States of the Former Soviet Union (CRDF), by the Russian Found of Basic Research, Grant No.00-02-17808 and INTAS Call 2000, Project No.587.

References

- [1] B.V.Geshkenbein, V.L.Morgunov, Phys.Lett. **B340** (1994) 185.
- [2] B.V.Geshkenbein, V.L.Morgunov, Phys.Lett. **B352** (1995) 456.
- [3] Particle Data Group, K.Hagiwara et al., Phys.Rev. **D66** (2000) 010001-1.
- [4] B.V.Geshkenbein, Yad.Fiz **59** (1996) 309.
- [5] B.V.Geshkenbein, hep-ph/0206094. Phys.Rev. **D67**, 0540XX (2003).
- [6] B.V.Geshkenbein, Yad.Fiz. **51** (1990) 1121.
- [7] B.V.Geshkenbein, V.L.Morgunov, Yad.Fiz. **58** (1995) 1873.
- [8] M.A.Shifman, A.I.Vainstein and V.I.Zakharov, Nucl.Phys. **B147** (1979) 385.
- [9] B.V.Geshkenbein, Yad.Fiz. **49** (1989) 1138.
- [10] M.Davier, S.Eidelman, A.Höcker and L.Zhang, hep-ph/0208177.
- [11] K.Hagiwara, A.D.Martin, Daisuke Nomura and T.Teubner, hep-ph/02091187 v2.
- [12] A.D.Martin, D.Zeppenfeld, Phys.Lett. **B345** (1995) 558.
- [13] S.Eidelman, F.Jegerlehner, Z.Phys. **C67** (1995) 585.
- [14] H.Burkhardt, B.Pietrzyk, Phys.Lett. **B356** (1995) 398.
- [15] M.L.Swartz, Phys.Rev. **D53** (1996) 5268.
- [16] M.Davier, A.Höcker, Phys.Lett. **B419** (1998) 419.
- [17] J.H.Kühn, M.Steinhauser, Phys.Lett. **B437** (1998) 425.
- [18] J.Erler, Phys.Rev. **D59** (1999) 054008.
- [19] M.Davier, A.Höcker, Phys.Lett. **B435** (1998) 427.
- [20] S.Groote et al., Phys.Lett. **B440** (1998) 375.
- [21] F.Jegerlehner, hep-ph/9901386.
- [22] A.D.Martin, J.Outhwaite, M.G.Ryskin, Phys.Lett. **B492** (2000) 69.